



ONDERSTEPSPOORT
BIOLOGICAL PRODUCTS

For animal use only

WESSELSBRON DISEASE VACCINE

Reg. No. G 0126 (Act 36/1947) Namibia: NSR 0954

Freeze-dried, live attenuated Wesselsbron virus for the immunisation of sheep and goats against Wesselsbron disease.

Store the vaccine in a refrigerator at a temperature of 4 °C to 8 °C. Do not use after the expiry date printed on the bottle.

RECOMMENDATIONS FOR USE

Animals can be vaccinated at any age except that lambs from immune animals should not be inoculated before they are six months old because maternal antibodies may block the vaccine response. This disease occurs during the late summer and autumn and it is, therefore, best to inoculate susceptible animals during spring, three to six weeks before the mating season. Where possible, breeding should also be arranged so that ewes lamb from February to April to ensure that lambs will be adequately protected by maternal antibodies during the time of the year that this disease occurs.

WARNINGS

Do not slaughter animals for human consumption within 7 days of vaccination. Vaccinate healthy animals only. Keep out of reach of children, uninformed persons and animals. Under no circumstances should pregnant animals be inoculated with this vaccine - as the virus can affect the fetus. Although this product has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of seasons. If this is suspected, seek veterinary advice and notify the registration holder.

DIRECTIONS FOR USE

Use only as directed.

Sterilise syringes and needles by boiling in water for at least 15 minutes. Do not use disinfectants or methylated spirits for sterilising either needles or syringes. The active ingredient of the vaccine is in the form of a powder or pellet in a small bottle. By means of a sterile syringe, transfer 5 ml diluent to the bottle containing the freeze-dried vaccine. Mix thoroughly until the powder is dissolved. Transfer this suspension back to the remaining diluent and again mix well. The vaccine is now ready for use and must be injected without delay. Avoid exposure to high temperatures and direct sunlight during inoculation. Shake the bottle before filling the syringe. Use a separate needle for each animal.

DOSAGE: Sheep and goats: 1 ml subcutaneously

EFFECTS OF THE VACCINE

A slight febrile reaction may occur on the second to fourth day following inoculation but subsides rapidly. Full immunity is obtained three weeks after inoculation. Immunity cannot be guaranteed in all animals.

PACKING

Available in bottles of 100 doses.

Registration holder:

Onderstepoort Biological Products SOC Ltd, Co. Reg. No. 2000/022686/06
Private Bag X07, Onderstepoort, 0110. Tel: +27 (0) 12 522 1500, Fax: +27 (0) 12 522 1591

Made in South Africa

Edition 3



WESSELSBRON-SIEKTE ENTSTOF

Reg. Nr. G 0126 (Wet 36/1947) Namibië: NSR 0954

Lewende, verswakte Wesselsbronsvirus in vriesgedroogde vorm vir die immunisering van skape en bokke teen Wesselsbronsiekte.

Bewaar die entstof in 'n yskas by 'n temperatuur van 4 °C tot 8 °C. Moet nie die entstof na die vervaldatum wat op die bottel gedruk is gebruik nie.

AANBEVELINGS VIR GEBRUIK

Diere kan op enige ouderdom ingeënt word behalwe dat lammers van gesoute diere nie ingeënt moet word voordat hulle ses maande oud is nie omdat die teenliggame in die biesmelk die uitwerking van die entstof mag belemmer. Hierdie siekte kom gedurende die laat somer en herfs voor en daarom is dit wenslik om vatbare diere gedurende die lente, 3–6 weke voordat die oöie by die ramme geplaas word, in te ent. Indien moontlik moet teling so gereël word dat oöie vanaf Februarie tot April lam om te verseker dat lammers beskerm is deur die teenliggame in die biesmelk gedurende dié tyd van die jaar wanneer hierdie siekte voorkom.

WAARSKUWINGS

Moet nie diere binne 7 dae na inenting vir menslike verbruik slag nie. Slegs gesonde diere moet ingeënt word. Hou buite bereik van kinders, oningeligte persone en diere. Dragtige diere moet onder geen omstandighede met hierdie entstof gespuit word nie -aangesien die entstof die fetus kan aantast. Alhoewel hierdie produk breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

GBRUIKSAANWYSINGS

Gebruik slegs soos voorgeskryf.

Steriliseer spuite en naalde deur dit vir ten minste 15 minute in water te kook. Moet nie ontsmettingsmiddels of brandspiritus gebruik om spuite en naalde te steriliseer nie. Die aktiewe bestanddeel van die entstof is in die vorm van 'n poeier of klont in 'n klein botteltjie. Dra 5 ml verdunningsvloeistof met 'n steriele spuit oor na die botteltjie gedroogde entstof. Meng deeglik totdat die poeier opgelos is. Dra die opgeloste entstof terug na die oorgeblewe verdunningsvloeistof en meng weer goed. Die entstof is nou gereed vir gebruik en moet dadelik ingespuit word. Vermoë blootstelling aan hoë temperature en direkte sonlig gedurende inenting. Skud die bottel goed voor gebruik. Gebruik 'n afsonderlike naald vir elke dier.

DOSIS: Skape en bokke: 1 ml onderhuids

UITWERKING VAN DIE ENTSTOF

'n Ligte koorsreaksie wat gou verdwyn mag vanaf die tweede tot vierde dag na inenting voorkom en na drie weke is die diere ten volle beskerm. Immuniteit kan nie in alle diere gewaarborg word nie.

VERPAKKING

Beskikbaar in bottels van 100 dosisse.

Registrasiehouer:

Onderstepoort Biologiese Produkte MSB Bpk, Mpy. Reg. Nr. 2000/022686/06
Privaatsak X07, Onderstepoort, 0110. Tel: +27 (0) 12 522 1500, Faks: +27 (0) 12 522 1591